**MIDAS™ MD1-59****MIDAS™:** a Modern Intelligent Dynamic Alternative Screen -

*A revolutionary 96 condition crystallization screen based on alternative polymeric precipitants<sup>1</sup>, developed and tested in the Laboratory of Dr. Clemens Grimm at University of Würzburg, Germany.*

MD1-59 is presented as a two box (96 x 10 mL conditions) kit.

**Features of MIDAS™**

- Ideal for both protein, protein/protein complexes, protein-nucleic acid complexes and sensitive macromolecular complexes.
- Narrow range of pH and salt concentrations centered on physiological values.
- Every condition contains at least one alternative polymeric precipitant.
- Designed to complement PEG and salt-based screens.
- Compatible with liquid-handling robots.

**Introduction:**

MIDAS is a 96 condition crystallization screen based on alternative polymeric precipitants. Devised and tested (Figure 1) in the Laboratory of Dr. Clemens Grimm *et al* of Würzburg University in Germany, MIDAS is a revolutionary crystallization screen that has moved away from the reliance on polyethylene glycols (PEGs) as the main precipitant (only 3 conditions in MIDAS contain a PEG).

For decades PEGs or their monomethyl ethers (PEG MMEs), have dominated crystallization screens. Out of 8289 entries scanned in the PDB, almost half of the crystallization conditions contained a PEG component and most commercial screens available today contain PEGs. However, the success rate of PEGs might be influenced due to their widespread dominance in crystallization screens.


**PEG Alternatives:**

There are many alternatives to PEGs and a variety have recently been described as being useful for macromolecular crystallogensis. Alternative polymers (Figure 2) such as the Jeffamine polyetheramines, pentaerythritol propoxylate and pentareytritol, polyvinyl pyrrolidone, polypropylene glycol, polyvinyl alcohol and polyacrylate have so far only sporadically been introduced into standard crystallization screens.

**MIDAS:**

To address this issue, recent work by Clemens Grimm *et al* have devised a protein crystallization screen (MIDAS) that systematically searches for crystallization conditions with alternative polymeric precipitants. To expand the precipitant diversity even more, they also scanned the current water-soluble polymer market for chemical variants or alternatives to the precipitants mentioned above. Owing to their particular properties as surface-active substances, ion exchangers and/or viscosity modifiers, many polymer variants have recently been developed.


MIDAS entails a relatively narrow range of pH and salt concentrations centered on physiological values to increase its suitability for sensitive macromolecular complexes, while every condition contains at least one alternative polymeric precipitant.



**Figure 1. Examples of protein crystals grown using MIDAS.**

(A) Lysozyme crystals obtained in 35% Sokalan HP 56, (B) Xylanase crystals obtained in 20% Jeffamine M2070, (C) Crystals of the cytokine receptor-ligand complex obtained in 45% pentaerythritol propoxylate (5/4 PO/OH).

(D) Crystals of streptavidin core obtained in 5% polyacrylate 2100, sodium salt, (E) Histone tail recognizing MBT repeats in 35% polyacrylate 2100, sodium salt, (F) Lysozyme crystals in 30% Sokalan CP 42, (G) spliceosomal assembly complex (SAC) 7 obtained in 6% polyvinyl pyrrolidone K15,

**Formulation Notes:**

MIDAS reagents are formulated using ultrapure water (>18.0 MΩ) and are sterile-filtered using 0.22 μm filters. No preservatives are added.

50% Stock solutions of Jeffamine are adjusted to pH 7.0 using HCl prior to inclusion in the reagents.

Sokalan® CP12 S was adjusted to pH 7 prior to using.

Final pH may vary from that specified on the datasheet. Molecular Dimensions will be happy to discuss the precise formulation of individual reagents. Individual reagents and stock solutions for optimization are available from Molecular Dimensions.

Enquiries regarding MIDAS formulation, interpretation of results or optimization strategies are welcome. Please e-mail, fax or phone your query to Molecular Dimensions.

Contact and product details can be found at [www.moleculardimensions.com](http://www.moleculardimensions.com)

Manufacturer's safety data sheets are available to download from our website.

**References :**

1. Grimm, C., Chari, A., Reuter, K. & Fischer, U. (2010). Acta Cryst. D66, 685-697.

**Ordering details:**

Catalogue Description	Catalogue Code
MIDAS™ 10 mL screen	MD1-59
MIDAS™ HT-96 screen	MD1-60
MIDAS™ 10 mL screen single reagents	MDSR-59-tube number
MIDAS™ HT-96 screen single reagents	MDSR-60-well number
For MIDAS™ stock reagents see our website under Optimization.	





Figure 2. Examples of alternative precipitants used in MIDAS™

\*R1 = —H for EO or —CH<sub>3</sub> for PO. The PO/EO molar ratio is 29/6 for Jeffamine M2005, 10/31 for Jeffamine M2070 and 9/1 for Jeffamine M600.

MIDAS™ is manufactured and distributed under an exclusive license with Dr. C. Grimm & Prof. Dr. U. Fischer.

Limited Use and Restrictions: Products sold by Molecular Dimensions Ltd. or its affiliates or authorized distributors and information relating to same are intended for research use only in crystal growth and optimization of crystal growth following use of the product by the purchaser and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, including resale or use in manufacture. The license to use MIDAS™ specifically excludes any rights to use the product information for the manufacture of the product or derivatives thereof, or distribute, transfer, or otherwise provide access to such information to any third party for any purpose or use.



Molecular  
Dimensions

moleculardimensions.com




MIDAS™

Box 1 of 2

MD1-59

Tube No.	% conc	Precipitant	% conc	Salt/Additive	pH	% conc	Buffer
1-1	50 % v/v	polypropylene glycol 400	5 %	dimethyl sulfoxide	6	0.1 M	HEPES-NaOH
1-2	12 % w/v	polyvinyl pyrrolidone K15	-	-	5.5	0.1 M	MES- NaOH
1-3	45 % w/v	polyacrylate 2100, sodium salt	-	-	6.5	0.1 M	HEPES-NaOH
1-4	14 % v/v	acrylic acid/maleic acid copolymer (50:50), sodium salt	-	-	-	-	-
1-5	12.5 % w/v	polyacrylate 2100, sodium salt	0.5 M	ammonium phosphate	-	-	-
1-6	19 % v/v	acrylic acid/maleic acid copolymer (50:50), sodium salt	-	-	8.5	0.1 M	Tris-HCl
1-7	10 % v/v	polypropylene glycol 400	-	-	-	-	-
1-8	5 % w/v	polyacrylate 2100, sodium salt	-	-	-	-	-
1-9	25 % v/v	pentaerythritol propoxylate (5/4 PO/OH)	-	-	6	0.1 M	MES- NaOH
1-10	24 % w/v	polyvinyl pyrrolidone K15	0.1 M	sodium sulfate	-	-	-
1-11	35 % v/v	pentaerythritol ethoxylate (15/4 EO/OH)	0.2 M	calcium chloride	6.5	0.1 M	HEPES-NaOH
1-12	35 % v/v	polypropylene glycol 400	-	-	7	0.1 M	K/Na Phosphate
1-13	20% v/v	Jeffamine D2000	0.2 M	sodium chloride	5.5	0.1 M	MES- NaOH
	10 % v/v	Jeffamine M2005	-	-	-	-	-
1-14	15 % v/v	pentaerythritol propoxylate (5/4 PO/OH)	0.2 M	sodium thiocyanate	7	0.1 M	HEPES-NaOH
1-15	5 % w/v	polyvinyl alcohol type II	0.2 M	potassium acetate	7	0.1 M	HEPES-NaOH
	10 % v/v	Jeffamine T403	-	-	-	-	-
1-16	45 % v/v	pentaerythritol propoxylate (5/4 PO/OH)	0.2 M	sodium chloride	6	0.1 M	MES- NaOH
1-17	8 % w/v	polyvinyl alcohol type II	10 % v/v	1- propanol	7	0.1 M	HEPES-NaOH
1-18	30 % w/v	polyvinyl pyrrolidone K15	0.1 M	lithium sulfate	7	0.1 M	HEPES-NaOH
1-19	40 % v/v	polypropylene glycol 400	0.2 M	imidazole	7	-	-
1-20	8 % w/v	acrylic acid/maleic acid copolymer (50:50), sodium salt	0.06 M	lithium sulfate	7.5	0.1 M	HEPES-NaOH
	3 % v/v	pentaerythritol ethoxylate (3/4 EO/OH)	-	-	-	-	-
1-21	35 % v/v	Jeffamine SD2001	0.1 M	sodium chloride	8	0.1 M	Tris-HCl
1-22	30 % v/v	Jeffamine M600	10 % v/v	dimethyl sulfoxide	-	-	-
1-23	20 % v/v	polypropylene glycol 400	10 % v/v	1-propanol	-	-	-
1-24	28 % w/v	acrylic acid/maleic acid copolymer (50:50), sodium salt	-	-	6.5	0.1 M	HEPES-NaOH
1-25	15 % w/v	Jeffamine ED2003	10 % v/v	ethanol	-	-	-
1-26	30 % w/v	Jeffamine ED2003	0.2 M	sodium chloride	6	0.1 M	MES- NaOH
1-27	25 % v/v	Jeffamine SD2001	0.1 M	sodium malonate	5.5	0.1 M	MES- NaOH
1-28	15 % v/v	pentaerythritol propoxylate (5/4 PO/OH)	0.2 M	sodium chloride	6	0.1 M	MES- NaOH
1-29	35 % v/v	pentaerythritol ethoxylate (3/4 EO/OH)	0.2 M	Magnesium chloride	-	-	-
1-30	40 % v/v	pentaerythritol propoxylate (5/4 PO/OH)	15 % v/v	ethanol	-	-	-
1-31	50 % v/v	pentaerythritol propoxylate (5/4 PO/OH)	-	-	8	0.1 M	Tris-HCl
1-32	12.5 % w/v	polyvinyl pyrrolidone K15	0.2 M	sodium chloride	8	0.1 M	Tris-HCl
	10 % w/v	PEG 4000	-	-	-	-	-
1-33	25 % v/v	pentaerythritol propoxylate (5/4 PO/OH)	0.1 M	sodium chloride	-	-	-
	10 % v/v	dimethyl sulfoxide,	-	-	-	-	-
1-34	35 % w/v	polyacrylate 2100, sodium salt	0.2 M	ammonium sulfate	7.5	0.1 M	HEPES-NaOH
1-35	30 % v/v	pentaerythritol ethoxylate (15/4 EO/OH)	0.1 M	magnesium formate	8.5	0.1 M	Tris-HCl
1-36	20 % v/v	Glascal W13	0.2 M	sodium sulfate	7.5	0.1 M	HEPES-NaOH
1-37	60 % v/v	polypropylene glycol 400	-	-	8	0.1 M	Tris-HCl
1-38	30 % v/v	pentaerythritol ethoxylate (15/4 EO/OH)	-	-	7.5	0.1 M	HEPES-NaOH
	6 % w/v	polyvinyl pyrrolidone K15	-	-	-	-	-
1-39	45 % v/v	polypropylene glycol 400	10 % v/v	ethanol	-	-	-
1-40	10 % v/v	pentaerythritol ethoxylate (3/4 EO/OH)	10 % v/v	1-butanol	-	-	-
1-41	12.5 % w/v	polyacrylate 2100, sodium salt	-	-	7	0.1 M	HEPES-NaOH
	6 % v/v	Jeffamine SD2001	-	-	-	-	-
1-42	6 % w/v	polyvinyl pyrrolidone K15	-	-	6.5	0.1 M	HEPES-NaOH
1-43	20 % w/v	Jeffamine ED2003	-	-	6.5	0.1 M	HEPES-NaOH
1-44	20 % v/v	glycerol ethoxylate	10 % v/v	tetrahydrofuran	8.0	0.1 M	Tris-HCl
1-45	25 % v/v	Jeffamine D2000	0.2 M	imidazole	7	-	-
1-46	30 % v/v	Jeffamine SD2001	0.2 M	potassium chloride	6.5	0.1 M	HEPES-NaOH
1-47	30 % v/v	polypropylene glycol 400	0.1 M	sodium chloride	-	-	-
1-48	20 % v/v	Jeffamine SD2001	15 % v/v	1-propanol	-	-	-

Recommended storage of kits is at room temperature (18°C) or 4°C. If stored at 4°C, turbidity of some solutions may occur. The effect is fully reversible; bringing the tubes back to room temperature and shaking results in clear solutions after 10 min and does not affect the quality of the solution.



MIDAS™

Box 2 of 2

MD1-59

Tube No.	% conc	Precipitant	% conc	Salt/Additive	pH	% conc	Buffer
2-1	25 % v/v	Jeffamine T403	0.2 M	lithium sulfate	8	0.1 M	Tris-HCl
2-2	35 % v/v	pentaerythritol propoxylate (5/4 EO/OH)	0.2 M	potassium acetate	-	-	-
2-3	20 % v/v	pentaerythritol ethoxylate (15/4 EO/OH)	0.2 M	potassium chloride	9.5	0.1 M	Glycine
2-4	40 % v/v	pentaerythritol propoxylate (5/4 PO/OH)	0.2 M	sodium thiocyanate	7	0.1 M	HEPES-NaOH
2-5	15 % v/v	Jeffamine T403	0.2 M	potassium chloride	6.5	0.1 M	HEPES-NaOH
	15 % v/v	Jeffamine ED2003	-	-	-	-	-
2-6	15 % v/v	pentaerythritol ethoxylate (15/4 EO/OH),	0.2 M	potassium acetate	6	0.1 M	MES- NaOH
	3 % v/v	Jeffamine T403	-	-	-	-	-
2-7	30 % w/v	polyacrylate 2100, sodium salt	0.1 M	sodium malonate	7	0.1 M	HEPES-NaOH
2-8	10 % v/v	Jeffamine D2000	10 % v/v	ethanol	-	-	-
	10 % v/v	Jeffamine M2005	-	-	-	-	-
2-9	25 % w/v	Jeffamine ED2003	0.1 M	lithium sulfate	8	0.1 M	Tris-HCl
2-10	10 % v/v	Jeffamine T403	-	-	8	0.1 M	Tris-HCl
	10 % w/v	Jeffamine ED2003	-	-	-	-	-
2-11	25 % w/v	polyacrylate 2100, sodium salt	0.1 M	lithium sulfate	6.5	0.1 M	HEPES-NaOH
2-12	15 % w/v	polyacrylate 2100, sodium salt	0.2 M	magnesium chloride	7.5	0.1 M	HEPES-NaOH
2-13	40 % v/v	Jeffamine D2000	-	-	6.5	0.1 M	HEPES-NaOH
2-14	10 % w/v	polyacrylate 2100, sodium salt	0.5 M	sodium chloride	8	0.1 M	Tris-HCl
2-15	14 % v/v	Jeffamine ED900	-	-	7	0.1 M	K/Na Phosphate
	11 % v/v	Jeffamine SD2001	-	-	-	-	-
2-16	20 % w/v	polyacrylate 2100, sodium salt	0.2 M	sodium chloride	9	0.1 M	Bicine
2-17	20 % v/v	Jeffamine D2000	0.2 M	sodium malonate	5.5	0.1 M	MES- NaOH
2-18	30 % v/v	Jeffamine M2070	0.2 M	potassium chloride	8	0.1 M	Tris-HCl
2-19	20 % v/v	Jeffamine M2070	20 % v/v	dimethyl sulfoxide	-	-	-
2-20	40 % w/v	pentaerythritol propoxylate (17/8 PO/OH)	0.2 M	magnesium chloride	5.5	0.1 M	MES- NaOH
2-21	20 % w/v	polyacrylate 5100, sodium salt	-	-	8	0.1 M	Tris-HCl
2-22	28 % v/v	poly(ethylene imine) branched	-	-	7	0.1 M	HEPES-NaOH
2-23	20 % v/v	Sokalan® CP 7	0.1 M	ammonium formate	7	0.1 M	HEPES-NaOH
2-24	20 % w/v	Sokalan® HP 56	0.2 M	sodium sulfate	8	0.1 M	Tris-HCl
2-25	25 % v/v	Sokalan® CP 7	0.1 M	potassium chloride	7	0.1 M	HEPES-NaOH
2-26	20 % v/v	Sokalan® CP 5	0.3 M	ammonium formate	7	0.1 M	HEPES-NaOH
2-27	40 % v/v	glycerol ethoxylate	-	-	-	-	-
2-28	30 % v/v	glycerol ethoxylate	-	-	8.5	0.1 M	Tris-HCl
2-29	15 % v/v	Sokalan® HP 66 K	-	-	7	0.1 M	HEPES-NaOH
	3 % v/v	poly(ethylene imine)	-	-	-	-	-
2-30	35 % v/v	glycerol ethoxylate	0.2 M	lithium citrate	-	-	-
2-31	30 % v/v	glycerol ethoxylate	0.2 M	ammonium acetate	6.5	0.1 M	MES- NaOH
2-32	20 % v/v	Sokalan® CP 42	5 % v/v	methanol	8	0.1 M	Tris-HCl
2-33	25 % v/v	Sokalan® CP 42	10 % v/v	tetrahydrofuran	7	0.1 M	Tris-HCl
2-34	20 % v/v	Sokalan® CP 42	0.1 M	lithium acetate	6	0.1 M	Bis-Tris- NaOH
2-25	15 % v/v	Sokalan® CP 12 S	0.1 M	lithium citrate	5.5	0.1 M	Bis-Tris- NaOH
2-36	15 % v/v	Sokalan® CP 5	-	-	6	0.1 M	Bis-Tris- NaOH
2-37	25 % v/v	Sokalan® CP 42	-	-	6	0.1 M	Bis-Tris- NaOH
2-38	25 % v/v	Sokalan® HP 66 K	0.2 M	ammonium acetate	7	0.1 M	HEPES-NaOH
2-39	20 % v/v	glycerol ethoxylate	-	-	8.5	0.1 M	Tris-HCl
	3 % v/v	poly(ethylene imine)	-	-	-	-	-
2-40	25 % v/v	glycerol ethoxylate	0.2 M	ammonium chloride	7.5	0.1 M	HEPES-NaOH
2-41	40 % v/v	Glasco® W13	0.2 M	potassium citrate	-	-	-
2-42	30 % w/v	polyacrylate 5100, sodium salt	10 % v/v	ethanol	6	0.1 M	MES- NaOH
2-43	15 % v/v	Sokalan® CP 42	0.2 M	potassium citrate	-	-	-
2-44	30 % v/v	Sokalan® CP 42	-	-	8.5	0.1 M	Tris-HCl
2-45	25 % w/v	Sokalan® HP 56	0.2 M	ammonium acetate	7	0.1 M	HEPES-NaOH
2-46	25 % v/v	Sokalan® CP 5	-	-	8.5	0.1 M	Tris-HCl
2-47	10 % w/v	poly(vinyl pyrrolidone) K15	0.2 M	ammonium formate	-	-	-
	20 % w/v	PEG 4000	-	-	-	-	-
2-48	15 % w/v	poly(vinyl pyrrolidone) K15	-	-	8	0.1 M	Tris-HCl
	25 % w/v	PEG MME 5000	-	-	-	-	-

Sokalan® are water-soluble polymers based on acrylic acid, maleic acid, vinylpyrrolidone, vinylimidazole and/or hydrophobic monomers.